

Client Name: _____ Date: _____

SPECIFIC SELF-MYOFASCIAL RELEASE TECHNIQUES



ILIOTIBIAL TRACT (IT Band)

Position yourself side lying on foam roll. Bottom leg is raised slightly off floor. Maintain head in "neutral" with ears aligned with shoulders. Roll just below hip joint down the lateral thigh to the knee.



PIRIFORMIS

Begin positioned as shown with foot crossed to opposite knee. Roll on the posterior hip area. Increase the stretch by pulling the knee toward the opposite shoulder



HAMSTRING

Place hamstrings on the roll with hips unsupported. Feet are crossed to increase leverage. Roll from knee toward posterior hip while keeping quadriceps tightened.



QUADRICEPS

Body is positioned prone with quadriceps on foam roll. It is very important to maintain proper Core control (abdominal Drawn-In position & tight gluteals) to prevent low back compensations. Roll from pelvic bone to knee, emphasizing the lateral thigh



TENSOR FASCIA LATAE (TFL)

Position the body as described above. Foam roll is placed just lateral to the anterior pelvic bone (ASIS). Roll slightly up and down along the upper portion of the front of your legs.



ADDUCTOR

Extend the thigh and place foam roll in the groin region with body prone on the floor. Be cautious when rolling near the adductor complex origins at the pelvis.



LATISSIMUS

Position yourself side lying with arm outstretched and foam roll placed in axillary (armpit) area. Thumb is pointed up to pre-stretch the ~~latissimus dorsi~~ muscle. Movement during this technique is minimal.



RHOMBOIDS

Cross arms to the opposite shoulder to clear the shoulder blades across the thoracic wall. While maintaining abdominal Draw-In position, raise hips until unsupported. Also stabilize the head in "neutral." Roll mid-back area on the foam.

NOTES: _____

Dr.Todd Plutchok, DC CCWP, ART Provider

The League Sports Injury Rehab
4510 Executive Drive, Plaza 5, San Diego CA 92121
Office (858) 452-3734

The Training Club
12264 El Camino Real #60 San Diego, CA 92130
Office (858) 793-0665 X 0


www.TheSportsAndWellnessDC.com email DrTodd@TheSportsAndWellnessDC.com

Client Name: _____ Date: _____

General Guidelines


1. Spend 1-2 minutes per self myofascial release technique and on each each side (when applicable).
2. When a trigger point is found (painful area) hold for 30-45 seconds.
3. Keep the abdominal muscles tight which provides stability to the lumbo-pelvic-hip complex during rolling.
4. Remember to breathe slowly as this will help to reduce any tense reflexes caused by discomfort.
5. Complete the self myofascial release exercises 1-2 x daily.

Muscle & Soft Tissue Treatments



Adhesive Tissue-Unorganized

Normal Tissue -Organized



Dr.Todd Plutchok, DC CCWP, ART Provider

The League Sports Injury Rehab
4510 Executive Drive, Plaza 5, San Diego CA 92121
Office (858) 452-3734

The Training Club
12264 El Camino Real #60 San Diego, CA 92130
Office (858) 793-0665 X 0

www.TheSportsAndWellnessDC.com email DrTodd@TheSportsAndWellnessDC.com